

# Cneorum

## Habitus and growth type

Plant height [m]: **0.8**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **9.05**

## Flower

Flowering period: **March-May, September-October**

## Fruit, seed and dispersal

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **9**

Moisture indicator value: **2.5**

Reaction indicator value: **5.5**

Nutrient indicator value: **3.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.37**

Disturbance frequency (herb layer): **1.5**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.3**

Mowing frequency: **0.05**

Grazing pressure: **0.29**

Soil disturbance: **0.14**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [FA \(QUI\) Quercetea ilicis](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [N1G Mediterranean coniferous coastal dune forest](#)

### Broad habitat

Occurrence in broad habitats: **Scrub, Forest**