Dodartia

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.06**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Disturbance indicator values

Disturbance frequency: **1.91**
Disturbance frequency (herb layer): **1.91**
Disturbance severity: **0.58**
Disturbance severity (herb layer): **0.57**
Mowing frequency: **0.53**
Grazing pressure: **0.27**
Soil disturbance: **0.26**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DB (FEP) *Festuco-Puccinellietea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R65 Continental subsaline alluvial pasture and meadow**, **S68 Semi-desert sand dune with sparse scrub**
Constant species of EUNIS habitats: **R65 Continental subsaline alluvial pasture and meadow**, **S68 Semi-desert sand dune with sparse scrub**

Broad habitat

Occurrence in broad habitats: **Saline vegetation, Semi-desert**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz