

# *Hispidella*

## Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **0.17**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Ellenberg-type indicator values

Light indicator value: **8.7**

Temperature indicator value: **7.3**

Moisture indicator value: **1.9**

Reaction indicator value: **3.9**

Nutrient indicator value: **2.1**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.64**

Disturbance frequency (herb layer): **2.06**

Disturbance severity: **0.31**

Disturbance severity (herb layer): **0.3**

Mowing frequency: **0.88**

Grazing pressure: **0.31**

Soil disturbance: **0.19**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FI \(TUB\) \*Helianthemetea guttati\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R1R Mediterranean to Atlantic open, dry, acid](#)

[and neutral grassland](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**