Chaiturus

Habitus and growth type

Plant height [m]: **0.84**
Life span: **Annual, Biennial or short-lived**
Life form: **Hemicryptophyte, Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **24.42**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.43**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**
Moisture indicator value: **5.8**
Reaction indicator value: **7.8**
Nutrient indicator value: **7.4**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **0.55**
Disturbance frequency (herb layer): **1.59**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.64**
Disturbance severity (herb layer): **0.48**
Mowing frequency: **0.3**
Grazing pressure: **0.18**
Soil disturbance: **0.2**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QF (ART) *Artemisietea vulgaris***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R65 Continental subsaline alluvial pasture and meadow**

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Saline vegetation, Scrub, Synanthropic**

Distribution

Continentality: **8**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz