Chamaedaphne

Habitus and growth type

Plant height [m]: **0.55**
Life span: **Perennial**
Life form: **Chamaephyte, Dwarf shrub**

Leaf

Specific leaf area [mm^2/mg]: **28.93**

Fruit, seed and dispersal

Seed mass [mg]: **0.09**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.8**
Temperature indicator value: **3.1**
Moisture indicator value: **8.3**
Reaction indicator value: **3**
Nutrient indicator value: **2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.25**
Disturbance frequency (herb layer): **1.37**
Disturbance severity: **0.53**
Disturbance severity (herb layer): **0.1**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.01**
Grazing pressure: **0.16**
Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **BA (PIC) *Vaccinio-Piceetea*, PB (OXY) *Oxycocco-Sphagnetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **Q11 Raised bog, T3J Pinus and Larix mire forest**
Constant species of EUNIS habitats: **T3J Pinus and Larix mire forest**

Broad habitat

Occurrence in broad habitats: **Mire, Forest**

Distribution

Continentality: **8**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz