

# *Chamaerops*

## Habitus and growth type

Plant height [m]: **2.63**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Spinescence: **spinescent**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **5.52**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]:

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **9.5**

Moisture indicator value: **3.5**

Reaction indicator value: **4.5**

Nutrient indicator value: **4.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.28**  
Disturbance frequency (herb layer): **1.4**  
Disturbance severity: **0.7**  
Disturbance severity (herb layer): **0.25**  
Mowing frequency: **0.02**  
Grazing pressure: **0.31**  
Soil disturbance: **0.13**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FA \(QUI\) Quercetea ilicis](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [S54 Thermomediterranean arid scrub](#), [S71 Western Mediterranean spiny heath](#)

Constant species of EUNIS habitats: [S54 Thermomediterranean arid scrub](#), [S71 Western Mediterranean spiny heath](#), [S72 Eastern Mediterranean spiny heath \(phrygana\)](#)

### Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**