

# *Myagrum*

## Habitus and growth type

Plant height [m]: **0.28**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **29.06**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Seed mass [mg]: **18.96**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **7.4**

Moisture indicator value: **3.9**

Reaction indicator value: **8.8**

Nutrient indicator value: **4.4**

Salinity indicator value: **0.2**

### Disturbance indicator values

Disturbance frequency: **2.09**

Disturbance frequency (herb layer): **2.09**

Disturbance severity: **0.82**

Disturbance severity (herb layer): **0.82**

Mowing frequency: **0.95**

Grazing pressure: **0.18**

Soil disturbance: **0.76**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [QA \(PAR\) \*Papaveretea rhoeadis\*](#)

### Broad habitat

Occurrence in broad habitats: **Synanthropic**

## **Distribution**

Continentality: 7

Continentality amplitude: 4