

# ***Radiola***

## **Habitus and growth type**

Plant height [m]: **0.06**

Life span: **Annual**

Life form: **Therophyte**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **32.29**

## **Flower**

Flowering period: **May-September**

## **Fruit, seed and dispersal**

Seed mass [mg]: **0.03**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Reaction indicator value: **3.8**

Nutrient indicator value: **2**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.74**

Disturbance frequency (herb layer): **1.92**

Disturbance severity: **0.61**

Disturbance severity (herb layer): **0.6**

Mowing frequency: **0.13**

Grazing pressure: **0.25**

Soil disturbance: **0.37**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [\*\*OC \(ISO\) Isoëto-Nanojuncetea\*\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [\*\*Q62 Periodically exposed shore with stable, mesotrophic sediments with pioneer or ephemeral vegetation\*\*](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland**

## Distribution

Continentality: **4**

Continentality amplitude: **4**