

Streptopus

Habitus and growth type

Plant height [m]: **0.6**

Life span: **Perennial**

Life form: **Geophyte**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **167.89**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **4.7**

Temperature indicator value: **3.6**

Reaction indicator value: **6.4**

Nutrient indicator value: **6.2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.08**

Disturbance frequency (herb layer): **1.27**

Disturbance severity: **0.65**

Disturbance severity (herb layer): **0.13**

Mowing frequency: **0**

Grazing pressure: **0.19**

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CR \(VIR\) *Betulo carpaticae-Alnetea viridis*](#), [CS \(MUL\) *Mulgedio-Aconitetea*](#)

Broad habitat

Occurrence in broad habitats: **Alpine-subalpine and arctic grassland, Scrub, Forest**

Distribution

Continentality amplitude: **5**