

# Taxus

## Habitus and growth type

Plant height [m]: **15.83**

Life span: **Perennial**

Life form: **Phanerophyte, Tree, Shrub**

## Leaf

Leaf type (woody plants only): **Needle-like**

Leaf phenology (woody plants only): **Evergreen**

Specific leaf area [mm<sup>2</sup>/mg]: **7.87**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Seed mass [mg]: **243.31**

Dispersal mode: **Endozoochory, Anthropochory**

Dispersal distance class: **6, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Reaction indicator value: **7.1**

Nutrient indicator value: **4.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.03**

Disturbance frequency (herb layer): **1.29**

Disturbance severity: **0.67**

Disturbance severity (herb layer): **0.13**

Mowing frequency: **0.01**

Grazing pressure: **0.19**

Soil disturbance: **0.11**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CA \(FAG\) \*Carpino-Fagetea sylvaticae\*](#),  
[CB \(PUB\) \*Quercetea pubescentis\*](#), [GC \(LAU\) \*Pruno lusitanicae-Lauretea azoricae\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T3C \*Taxus baccata\* forest](#)

Constant species of EUNIS habitats: [T3C \*Taxus baccata\* forest](#)

Dominant species of EUNIS habitats: [T3C \*Taxus baccata\* forest](#)

### Broad habitat

Occurrence in broad habitats: **Forest, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **4**

Continentality amplitude: **4**