

Leontodon crispus aggr.

Habitus and growth type

Plant height [m]: **0.28**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **2.33**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **2.27**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **6.3**

Moisture indicator value: **3.4**

Reaction indicator value: **7.7**

Nutrient indicator value: **1.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.97**

Disturbance frequency (herb layer): **1.79**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.4**

Mowing frequency: **0.34**

Grazing pressure: **0.35**

Soil disturbance: **0.18**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#)

Constant species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#),
[R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region](#), [S34 Balkan-Anatolian submontane genistoid scrub](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Oromediterranean scrub, Scrub, Sparsely vegetated (incl. rock and scree)**