

# *Santolina chamaecyparissus* aggr.

## Habitus and growth type

Plant height [m]: **0.31**

Life span: **Perennial**

Life form: **Chamaephyte**

Spinescence: **not spinescent**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **8.92**

## Flower

Flowering period: **March-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.35**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic, Slightly acidic to near-neutral, Alkaline**

Nutrient relationship: **Dystrophic, Oligotrophic, Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.8**

Temperature indicator value: **8.5**

Moisture indicator value: **2.3**

Nutrient indicator value: **4.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.87**

Disturbance frequency (herb layer): **1.54**

Disturbance severity: **0.58**

Disturbance severity (herb layer): **0.39**

Mowing frequency: **0.15**

Grazing pressure: **0.35**

Soil disturbance: **0.22**

## **Habitat and sociology**

EUNIS habitat

Diagnostic species of EUNIS habitats: [S61 Western basiphilous garrigue](#), [S65 Mediterranean gypsum scrub](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**