Sparganium erectum aggr.

Habitus and growth type

Plant height [m]: **1.05**
Life span: **Perennial**
Life form: **Hydrophyte**

Leaf

Specific leaf area [mm^2/mg]: **19.51**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **17.21**
Dispersal mode: **Endozoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.2**
Moisture indicator value: **10.1**
Reaction indicator value: **6.7**
Nutrient indicator value: **6.2**
Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **1.45**
Disturbance frequency (herb layer): **1.74**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.28**
Disturbance severity (herb layer): **0.26**
Mowing frequency: **0.24**
Grazing pressure: **0.09**
Soil disturbance: **0.07**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **P3c Fresh-water large pleustophyte vegetation**, **Q51 Tall-helophyte bed**
Constant species of EUNIS habitats: **P3c Fresh-water large pleustophyte vegetation**, **Q51 Tall-helophyte bed**
Dominant species of EUNIS habitats: **Q51 Tall-helophyte bed**

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz