

# *Anacamptis palustris* aggr.

## Habitus and growth type

Plant height [m]: **0.4**

Life span: **Perennial**

Life form: **Geophyte**

## Flower

Flowering period: **March-July**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **initial or partial mycoheterotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **8.3**

Moisture indicator value: **8.1**

Reaction indicator value: **7.7**

Salinity indicator value: **0.9**

### Disturbance indicator values

Disturbance frequency: **1.46**

Disturbance frequency (herb layer): **2.04**

Disturbance severity: **0.47**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.65**

Grazing pressure: **0.27**

Soil disturbance: **0.15**

## Habitat and sociology

### EUNIS habitat

Diagnostic species of EUNIS habitats: [N1J Mediterranean and Black Sea moist and wet dune slack](#), [R34 Submediterranean moist meadow](#)

Constant species of EUNIS habitats: [R34 Submediterranean moist meadow](#)

### Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Aquatic, Wetland, Mire, Grassland (non-alpine, non-saline), Saline vegetation, Forest**