Anacamptis palustris aggr.

Habitus and growth type

Plant height [m]: **0.4**
Life span: **Perennial**
Life form: **Geophyte**

Flower

Flowering period: **March-July**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **initial or partial mycoheterotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8.3**
Moisture indicator value: **8.1**
Reaction indicator value: **7.7**
Salinity indicator value: **0.9**

Disturbance indicator values

Disturbance frequency: **1.46**
Disturbance frequency (herb layer): **2.04**
Disturbance severity: **0.47**
Disturbance severity (herb layer): **0.45**
Mowing frequency: **0.65**
Grazing pressure: **0.27**
Soil disturbance: **0.15**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **N1J Mediterranean and Black Sea moist and wet dune slack**, **R34 Submediterranean moist meadow**

Constant species of EUNIS habitats: **R34 Submediterranean moist meadow**

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Aquatic, Wetland, Mire, Grassland (non-alpine, non-saline), Saline vegetation, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz