

# *Crepis neglecta* aggr.

## Habitus and growth type

Plant height [m]: **0.29**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **March-July**

## Fruit, seed and dispersal

Seed mass [mg]: **0.09**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **7.5**

Moisture indicator value: **3.5**

Reaction indicator value: **7**

Nutrient indicator value: **6**

Salinity indicator value: **0.5**

### Disturbance indicator values

Disturbance frequency: **0.9**

Disturbance frequency (herb layer): **1.86**

Disturbance severity: **0.75**

Disturbance severity (herb layer): **0.58**

Mowing frequency: **0.22**

Grazing pressure: **0.35**

Soil disturbance: **0.32**

## **Habitat and sociology**

EUNIS habitat

Diagnostic species of EUNIS habitats: [S63 Eastern garrigue](#), [S72 Eastern Mediterranean spiny heath \(phrygana\)](#)

Constant species of EUNIS habitats: [S63 Eastern garrigue](#), [S72 Eastern Mediterranean spiny heath \(phrygana\)](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**