

Dryopteris carthusiana aggr.

Habitus and growth type

Plant height [m]: **0.7**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **22.19**

Flower

Flowering period: **April-November**

Fruit, seed and dispersal

Seed mass [mg]: **0**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Reaction indicator value: **3.6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.09**

Disturbance frequency (herb layer): **1.22**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.16**

Mowing frequency: **0.01**

Grazing pressure: **0.19**

Soil disturbance: **0.13**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [T15 Broadleaved swamp forest on non-acid peat](#), [T31 Temperate mountain Picea forest](#)

Constant species of EUNIS habitats: [N1D Atlantic and Baltic broad-leaved coastal dune forest](#), [N1F Baltic coniferous coastal dune forest](#), [S25 Subalpine and subarctic deciduous scrub](#), [S26 Subalpine Pinus mugo scrub](#), [S32 Temperate Rubus scrub](#), [S37 Corylus avellana scrub](#), [S38 Temperate forest clearing scrub](#), [S92 Salix fen scrub](#), [T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils](#), [T13 Temperate hardwood riparian forest](#), [T15 Broadleaved swamp forest on non-acid peat](#), [T16 Broadleaved mire forest on acid peat](#), [T17 Fagus forest on non-acid soils](#), [T18 Fagus forest on acid soils](#), [T1B Acidophilous Quercus forest](#), [T1E Carpinus and Quercus mesic deciduous forest](#), [T1F Ravine forest](#), [T1H Broadleaved deciduous plantation of non site-native trees](#), [T27 Ilex aquifolium forest](#), [T31 Temperate mountain Picea forest](#), [T32 Temperate mountain Abies forest](#), [T34 Temperate subalpine Larix, Pinus cembra and Pinus uncinata forest](#), [T35 Temperate continental Pinus sylvestris forest](#), [T3F Dark taiga](#), [T3G Pinus sylvestris light taiga](#), [T3K Picea mire forest](#), [T3M Coniferous plantation of non site-native trees](#), [U23 Temperate, lowland to montane siliceous scree](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**