

Dryopteris filix-mas aggr.

Habitus and growth type

Plant height [m]: **0.75**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **24.2**

Flower

Flowering period: **May-October**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **3**

Temperature indicator value: **5**

Moisture indicator value: **5**

Reaction indicator value: **5**

Nutrient indicator value: **6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.05**

Disturbance frequency (herb layer): **1.29**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.15**

Mowing frequency: **0.01**

Grazing pressure: **0.19**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **BB (ASA) *Asaro europaei-Abietetea sibiricae*, CA (FAG) *Carpino-Fagetea sylvaticae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **[T17 Fagus forest on non-acid soils](#), [T1F Ravine forest](#), [T31 Temperate mountain Picea forest](#), [T32 Temperate mountain Abies forest](#)**

Constant species of EUNIS habitats: **[R56 Montane to subalpine moist or wet tall-herb and fern fringe](#), [S25 Subalpine and subarctic deciduous scrub](#), [S37 Corylus avellana scrub](#), [S38 Temperate forest clearing scrub](#), [T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils](#), [T17 Fagus forest on non-acid soils](#), [T1D Southern European mountain Betula and Populus tremula forest on mineral soils](#), [T1E Carpinus and Quercus mesic deciduous forest](#), [T1F Ravine forest](#), [T27 Ilex aquifolium forest](#), [T31 Temperate mountain Picea forest](#), [T32 Temperate mountain Abies forest](#), [T39 Mediterranean and Balkan subalpine Pinus heldreichii-Pinus peuce forest](#), [T3C Taxus baccata forest](#)**

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **5**