

Galium palustre aggr.

Habitus and growth type

Plant height [m]: **0.7**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **34.3**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **1.02**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.7**

Temperature indicator value: **5.1**

Moisture indicator value: **8.9**

Reaction indicator value: **5.2**

Nutrient indicator value: **4.3**

Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **0.89**

Disturbance frequency (herb layer): **1.63**
Disturbance severity: **0.44**
Disturbance severity (herb layer): **0.36**
Mowing frequency: **0.4**
Grazing pressure: **0.21**
Soil disturbance: **0.13**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q53 Tall-sedge bed](#), [R35 Moist or wet mesotrophic to eutrophic hay meadow](#), [S92 Salix fen scrub](#), [T15 Broadleaved swamp forest on non-acid peat](#)

Constant species of EUNIS habitats: [MA232 Baltic coastal meadow](#), [N1H Atlantic and Baltic moist and wet dune slack](#), [Q24 Intermediate fen and soft-water spring mire](#), [Q42 Extremely rich moss-sedge fen](#), [Q51 Tall-helophyte bed](#), [Q52 Small-helophyte bed](#), [Q53 Tall-sedge bed](#), [Q54 Inland saline or brackish helophyte bed](#), [R34 Submediterranean moist meadow](#), [R35 Moist or wet mesotrophic to eutrophic hay meadow](#), [R36 Moist or wet mesotrophic to eutrophic pasture](#), [R37 Temperate and boreal moist or wet oligotrophic grassland](#), [S92 Salix fen scrub](#), [T11 Temperate Salix and Populus riparian forest](#), [T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils](#), [T13 Temperate hardwood riparian forest](#), [T15 Broadleaved swamp forest on non-acid peat](#), [T16 Broadleaved mire forest on acid peat](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Aquatic, Spring, Wetland, Mire, Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**