

# *Melampyrum nemorosum* aggr.

## Habitus and growth type

Plant height [m]: **0.4**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **38.18**

## Flower

Flowering period: **May-August**

## Fruit, seed and dispersal

Seed mass [mg]: **7.94**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **5.1**

Moisture indicator value: **4.6**

Reaction indicator value: **5.4**

Nutrient indicator value: **3.8**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.13**

Disturbance frequency (herb layer): **1.38**

Disturbance severity: **0.65**

Disturbance severity (herb layer): **0.19**

Mowing frequency: **0.09**

Grazing pressure: **0.2**

Soil disturbance: **0.12**

## **Habitat and sociology**

EUNIS habitat

Diagnostic species of EUNIS habitats: [R51 Thermophilous forest fringe of base-rich soils](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**