

Ranunculus montanus aggr.

Habitus and growth type

Plant height [m]: **0.22**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **1.55**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **3.5**

Moisture indicator value: **4.8**

Reaction indicator value: **6.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.54**

Disturbance frequency (herb layer): **1.24**

Disturbance severity: **0.44**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.11**

Grazing pressure: **0.26**

Soil disturbance: **0.13**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R44 Arctic-alpine calcareous grassland](#), [T34 Temperate subalpine Larix, Pinus cembra and Pinus uncinata forest](#)

Constant species of EUNIS habitats: [R23 Mountain hay meadow](#), [R41 Snow-bed vegetation](#), [R43 Temperate acidophilous alpine grassland](#), [R44 Arctic-alpine calcareous grassland](#), [R45 Alpine and subalpine calcareous grassland of the Balkans and Apennines](#), [S21 Subarctic and alpine dwarf Salix scrub](#), [S26 Subalpine Pinus mugo scrub](#), [T34 Temperate subalpine Larix, Pinus cembra and Pinus uncinata forest](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**