Ribes rubrum aggr.

Habitus and growth type

Plant height [m]: **1.25**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**
Leaf phenology (woody plants only): **Winter deciduous**
Specific leaf area [mm^2/mg]: **31.19**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **74.42**
Dispersal mode: **Endozoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **4.4**
Temperature indicator value: **5**
Moisture indicator value: **7.4**
Reaction indicator value: **6.3**
Nutrient indicator value: **6.1**
Salinity indicator value: **0**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance indicator values

Disturbance frequency: **0.09**
Disturbance frequency (herb layer): **1.42**
Disturbance severity: **0.7**
Disturbance severity (herb layer): **0.24**
Mowing frequency: **0.01**
Grazing pressure: **0.19**
Soil disturbance: **0.12**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **N1D Atlantic and Baltic broad-leaved coastal dune forest**, **T13 Temperate hardwood riparian forest**
Constant species of EUNIS habitats: **N1D Atlantic and Baltic broad-leaved coastal dune forest**, **T13 Temperate hardwood riparian forest**

Broad habitat

Occurrence in broad habitats: **Alpine-subalpine and arctic grassland, Scrub, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz