

Sideritis hyssopifolia aggr.

Habitus and growth type

Plant height [m]: **0.25**

Life span: **Perennial**

Life form: **Chamaephyte**

Flower

Flowering period: **June-September**

Fruit, seed and dispersal

Seed mass [mg]: **1.39**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.7**

Moisture indicator value: **2.6**

Reaction indicator value: **6.8**

Nutrient indicator value: **1.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.23**

Disturbance frequency (herb layer): **1.79**

Disturbance severity: **0.45**

Disturbance severity (herb layer): **0.38**

Mowing frequency: **0.27**

Grazing pressure: **0.34**

Soil disturbance: **0.15**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [**R1H Iberian oromediterranean basiphilous dry grassland**](#)

Constant species of EUNIS habitats: [**R1H Iberian oromediterranean basiphilous dry grassland**](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**