

***Thymus odoratissimus* aggr.**

Habitus and growth type

Plant height [m]: **0.15**

Life span: **Perennial**

Life form: **Chamaephyte**

Leaf

Leaf type (woody plants only): **Sclerophyllous**

Leaf phenology (woody plants only): **Evergreen**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.14**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**

Moisture indicator value: **3.1**

Nutrient indicator value: **2.1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.35**

Disturbance frequency (herb layer): **1.89**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.58**

Grazing pressure: **0.33**

Soil disturbance: **0.17**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1B Continental dry grassland \(true steppe\)](#),
[S34 Balkan-Anatolian submontane genistoid scrub](#), [S36 Low steppic scrub](#)

Constant species of EUNIS habitats: [R1B Continental dry grassland \(true steppe\)](#),
[S34 Balkan-Anatolian submontane genistoid scrub](#), [S36 Low steppic scrub](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**