

Taraxacum sect. Alpina

Habitus and growth type

Plant height [m]: **0.13**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **27.4**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic, Wet**

Substrate reaction relationship: **Acidic, Alkaline**

Nutrient relationship: **Dystrophic, Oligotrophic, Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **2.1**

Moisture indicator value: **6.3**

Reaction indicator value: **6.5**

Nutrient indicator value: **5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.56**

Disturbance frequency (herb layer): **1.59**

Disturbance severity: **0.27**

Disturbance severity (herb layer): **0.27**

Mowing frequency: **0.06**

Grazing pressure: **0.22**

Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KE \(THL\) *Thlaspietea rotundifolii*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q23 Relict mire of Mediterranean mountains](#),
[R41 Snow-bed vegetation](#)

Constant species of EUNIS habitats: [Q23 Relict mire of Mediterranean mountains](#),
[R41 Snow-bed vegetation](#)

Broad habitat

Occurrence in broad habitats: **Spring, Mire, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **3**

Continentality amplitude: **3**