

Taraxacum sect. Erythrosperma

Habitus and growth type

Plant height [m]: **0.13**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **20.22**

Flower

Flowering period: **February-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.41**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.16**

Disturbance frequency (herb layer): **1.62**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.42**

Mowing frequency: **0.34**

Grazing pressure: **0.32**

Soil disturbance: **0.26**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1K Balkan and Anatolian oromediterranean dry grassland](#)

Constant species of EUNIS habitats: [N1A Atlantic and Baltic coastal dune scrub](#), [R13 Cryptogam- and annual-dominated vegetation on calcareous and ultramafic rock outcrops](#), [R1K Balkan and Anatolian oromediterranean dry grassland](#)

Broad habitat

Occurrence in broad habitats: **Spring, Grassland (non-alpine, non-saline), Oromediterranean scrub, Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **6**