Taraxacum sect. Obliqua

Habitus and growth type

Plant height [m]: **0.07**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **16.63**

Flower

Flowering period: **February-August**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry, Mesic**
Substrate reaction relationship: **Acidic, Slightly acidic to near-neutral**
Nutrient relationship: **Dystrophic, Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **3.5**
Moisture indicator value: **3**
Reaction indicator value: **6**
Nutrient indicator value: **4**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.72**
Disturbance frequency (herb layer): **1.76**
Disturbance severity: **0.35**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.34**
Mowing frequency: **0.16**
Grazing pressure: **0.32**
Soil disturbance: **0.18**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **4**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz