Taraxacum sect. Obovata

Habitus and growth type

Plant height [m]: **0.08**
Life form: **Hemicryptophyte**

Flower

Flowering period: **April-July**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**
Temperature indicator value: **7.5**
Moisture indicator value: **3**
Reaction indicator value: **6.3**
Nutrient indicator value: **6.3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.86**
Disturbance frequency (herb layer): **1.58**
Disturbance severity: **0.7**
Disturbance severity (herb layer): **0.57**
Mowing frequency: **0.43**
Grazing pressure: **0.3**
Soil disturbance: **0.25**

Habitat and sociology

Broad habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz