Acer campestre

Habitus and growth type

Plant height [m]: **13.92**
Life span: **Perennial**
Life form: **Phanerophyte, Tree**

Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**
Leaf phenology (woody plants only): **Winter deciduous**
Specific leaf area [mm^2/mg]: **18.93**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **73.61**
Dispersal mode: **Anemochory, Anthropochory**
Dispersal distance class: **4, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.7**
Reaction indicator value: **7.3**
Nutrient indicator value: **5.4**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.04**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.32**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.14**
Mowing frequency: **0.01**
Grazing pressure: **0.2**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CA (FAG) *Carpino-Fagetea sylvaticae***, **CD (RHA) *Crataego-Prunetea***, **HA (POP) *Alno glutinosae-Populetea albae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T19 Temperate and submediterranean thermophilous deciduous forest**, **T1E Carpinus and Quercus mesic deciduous forest**, **T1F Ravine forest**, **T3C Taxus baccata forest**
Constant species of EUNIS habitats: **S37 Corylus avellana scrub**, **T13 Temperate hardwood riparian forest**, **T17 Fagus forest on non-acid soils**, **T19 Temperate and submediterranean thermophilous deciduous forest**, **T1E Carpinus and Quercus mesic deciduous forest**, **T1F Ravine forest**, **T1H Broadleaved deciduous plantation of non site-native trees**, **T3C Taxus baccata forest**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality amplitude: **6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz