

# *Aconogonon alpinum*

## Habitus and growth type

Plant height [m]:

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **June-August**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **3.3**

Moisture indicator value: **4.8**

Reaction indicator value: **4.5**

Nutrient indicator value: **3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.57**

Disturbance frequency (herb layer): **1.54**

Disturbance severity: **0.49**

Disturbance severity (herb layer): **0.33**

Mowing frequency: **0.22**

Grazing pressure: **0.24**

Soil disturbance: **0.12**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) \*Molinio-Arrhenatheretea\*](#), [CR \(VIR\) \*Betulo carpaticae-Alnetea viridis\*](#), [KE \(THL\) \*Thlaspietea rotundifolii\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T3H \*Larix light taiga\*](#)

Constant species of EUNIS habitats: [T3H \*Larix light taiga\*](#)

### Broad habitat

Occurrence in broad habitats: **Spring, Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: 7

Continentality amplitude: 5