# *Adoxa moschatellina*

### Habitus and growth type

Plant height [m]: **0.1**
Life span: **Perennial**
Life form: **Hemicryptophyte, Geophyte**

### Leaf

Specific leaf area [$mm^2$/mg]: **38.45**

### Flower

Flowering period: **February-May**

### Fruit, seed and dispersal

Seed mass [mg]: **38.21**
Dispersal mode: **Endozoochory**
Dispersal distance class: **6**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **Native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

#### Ellenberg-type indicator values

Light indicator value: **4.3**
Temperature indicator value: **4.8**
Moisture indicator value: **6.1**
Reaction indicator value: **6.9**
Nutrient indicator value: **7.3**
Salinity indicator value: **0**

#### Disturbance indicator values

Disturbance frequency: **0.08**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Disturbance frequency (herb layer): **1.32**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.19**
Mowing frequency: **0.01**
Grazing pressure: **0.18**
Soil disturbance: **0.1**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **BB (ASA) *Asaro europaei-Abietetea sibiricae*, CA (FAG) *Carpino-Fagetea sylvaticae***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **T1F Ravine forest**
Constant species of EUNIS habitats: **T1F Ravine forest**

### Broad habitat

Occurrence in broad habitats: **Spring, Scrub, Forest, Synanthropic**

## Distribution

Continentality: **6**
Continentality amplitude: **5**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*