

# *Aegonychon purpureocaeruleum*

## Habitus and growth type

Plant height [m]: **0.41**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **37.01**

## Flower

Flowering period: **March-June**

## Fruit, seed and dispersal

Seed mass [mg]: **14.85**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **5.4**

Moisture indicator value: **3.9**

Reaction indicator value: **7.7**

Nutrient indicator value: **3.9**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.03**

Disturbance frequency (herb layer): **1.31**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.12**

Mowing frequency: **0.02**

Grazing pressure: **0.21**

Soil disturbance: **0.1**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) \*Quercetea pubescentis\*](#), [CL \(GER\) \*Trifolio-Geranietea sanguinei\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T19 Temperate and submediterranean thermophilous deciduous forest](#)

Constant species of EUNIS habitats: [T19 Temperate and submediterranean thermophilous deciduous forest](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

## Distribution

Continentality: **6**

Continentality amplitude: **5**