

# *Agropyron dasyanthum*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **5.5**

Moisture indicator value: **5**

Disturbance indicator values

Disturbance frequency: **1.63**

Disturbance frequency (herb layer): **1.66**

Disturbance severity: **0.36**

Disturbance severity (herb layer): **0.36**

Mowing frequency: **0.28**

Grazing pressure: **0.13**

Soil disturbance: **0.42**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CJ \(COR\) \*Koelerio-Corynephoretea canescentis\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R11 Pannonian and Pontic sandy steppe](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**