

# *Agrostis rupestris*

## Habitus and growth type

Plant height [m]: **0.08**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **8.53**

## Flower

Flowering period: **June-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.11**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.9**

Temperature indicator value: **1.8**

Moisture indicator value: **4.3**

Reaction indicator value: **2.2**

Nutrient indicator value: **1**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.4**

Disturbance frequency (herb layer): **1.5**

Disturbance severity: **0.29**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.01**

Grazing pressure: **0.25**

Soil disturbance: **0.09**

## Habitat and sociology

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R43 Temperate acidophilous alpine grassland](#)

Constant species of EUNIS habitats: [R1J Cyrno-Sardean oromediterranean siliceous dry grassland](#), [R43 Temperate acidophilous alpine grassland](#), [S21 Subarctic and alpine dwarf Salix scrub](#), [S22 Alpine and subalpine ericoid heath](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Heathland, Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **5**

Continentality amplitude: **3**