

Agrostis vinealis

Habitus and growth type

Plant height [m]: **0.27**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **19.87**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.07**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Reaction indicator value: **2.6**

Nutrient indicator value: **1.6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1**

Disturbance frequency (herb layer): **1.72**

Disturbance severity: **0.45**

Disturbance severity (herb layer): **0.36**

Mowing frequency: **0.27**

Grazing pressure: **0.3**

Soil disturbance: **0.25**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CJ \(COR\) *Koelerio-Coryneporetea canescentis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1Q Inland sanddrift and dune with siliceous grassland](#)

Constant species of EUNIS habitats: [R1P Oceanic to subcontinental inland sand grassland on dry acid and neutral soils](#), [R1Q Inland sanddrift and dune with siliceous grassland](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality amplitude: **8**