

Achillea atrata

Habitus and growth type

Plant height [m]: **0.13**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.34**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **1.5**

Moisture indicator value: **5**

Reaction indicator value: **8**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.37**

Disturbance frequency (herb layer): **1.4**

Disturbance severity: **0.29**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.02**

Grazing pressure: **0.15**

Soil disturbance: **0.17**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KE \(THL\) *Thlaspietea rotundifolii*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R41 Snow-bed vegetation](#), [U26 Temperate high-mountain base-rich scree and moraine](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Alpine-subalpine and arctic grassland, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **5**

Continentality amplitude: **1**