Achillea micrantha

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **5.8**
Moisture indicator value: **4**

Disturbance indicator values

Disturbance frequency: **1.67**
Disturbance frequency (herb layer): **1.79**
Disturbance severity: **0.41**
Disturbance severity (herb layer): **0.41**
Mowing frequency: **0.35**
Grazing pressure: **0.22**
Soil disturbance: **0.39**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **BD (PYR) *Pyrolo-Pinetea sylvestris***, **CJ (COR) *Koelerio-Corynephoretea canescentis***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R11 Pannonian and Pontic sandy steppe**, **S68 Semi-desert sand dune with sparse scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Constant species of EUNIS habitats: **R11 Pannonian and Pontic sandy steppe**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz