Aira cupaniana

Habitus and growth type

Plant height [m]: **0.19**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **0.05**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Dystrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.5**
Temperature indicator value: **9**
Moisture indicator value: **2**
Reaction indicator value: **3**
Nutrient indicator value: **1**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.28**
Disturbance frequency (herb layer): **1.95**
Disturbance severity: **0.51**
Disturbance severity (herb layer): **0.47**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.47**
Grazing pressure: **0.3**
Soil disturbance: **0.27**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FI (TUB) *Helianthemetea guttati***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1R Mediterranean to Atlantic open, dry, acid and neutral grassland**
Constant species of EUNIS habitats: **R1R Mediterranean to Atlantic open, dry, acid and neutral grassland**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz