Allium frigidum

Habitus and growth type

Life span: **Perennial**
Life form: **Geophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.1**
Temperature indicator value: **7.5**
Moisture indicator value: **3.8**
Reaction indicator value: **7.1**
Nutrient indicator value: **4**
Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **2.15**
Disturbance frequency (herb layer): **2.15**
Disturbance severity: **0.37**
Disturbance severity (herb layer): **0.32**
Mowing frequency: **0**
Grazing pressure: **0.36**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FR (DAP) *Daphno-Festucetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1K Balkan and Anatolian oromediterranean dry grassland**, **S75 Eastern Mediterranean mountain hedgehog-heath**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Broad habitat

Occurrence in broad habitats: **Oromediterranean scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz