

# *Allium victorialis*

## Habitus and growth type

Plant height [m]: **0.45**

Life span: **Perennial**

Life form: **Geophyte**

## Flower

Flowering period: **June-August**

## Fruit, seed and dispersal

Seed mass [mg]: **6.43**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Environmental relationships

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.38**

Disturbance frequency (herb layer): **0.96**

Disturbance severity: **0.47**

Disturbance severity (herb layer): **0.23**

Mowing frequency: **0.04**

Grazing pressure: **0.21**

Soil disturbance: **0.09**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CS \(MUL\) Mulgedio-Aconitetea](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Sparsely vegetated (incl. rock and scree)**

### **Distribution**

Continentality: **4**

Continentality amplitude: **4**