

Allium vineale

Habitus and growth type

Plant height [m]: **0.5**

Life span: **Perennial**

Life form: **Geophyte**

Leaf

Specific leaf area [mm²/mg]: **18.7**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **5.21**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.4**

Temperature indicator value: **6.4**

Moisture indicator value: **4.1**

Reaction indicator value: **6.4**

Nutrient indicator value: **6.4**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.04**

Disturbance frequency (herb layer): **1.87**
Disturbance severity: **0.56**
Disturbance severity (herb layer): **0.5**
Mowing frequency: **0.67**
Grazing pressure: **0.2**
Soil disturbance: **0.35**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [QA \(PAR\) Papaveretea rhoeadis](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **4**
Continentality amplitude: **4**