

# *Allosorus pteridioides*

## Habitus and growth type

Plant height [m]: **0.12**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **February-May**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.7**

Moisture indicator value: **2.8**

Reaction indicator value: **5**

Nutrient indicator value: **2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.24**

Disturbance frequency (herb layer): **1.56**

Disturbance severity: **0.64**

Disturbance severity (herb layer): **0.59**

Mowing frequency: **0.75**

Grazing pressure: **0.39**

Soil disturbance: **0.19**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [KC \(ASP\) \*Asplenietea trichomanis\*](#)

### Broad habitat

Occurrence in broad habitats: **Coastal cliff, Sparsely vegetated (incl. rock and scree)**