

# *Alopecurus rendlei*

## Habitus and growth type

Plant height [m]: **0.33**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **March-June**

## Fruit, seed and dispersal

Seed mass [mg]: **2.66**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.7**

Temperature indicator value: **8.3**

Moisture indicator value: **7.7**

Reaction indicator value: **7**

Nutrient indicator value: **7**

Salinity indicator value: **0.7**

### Disturbance indicator values

Disturbance frequency: **1.79**

Disturbance frequency (herb layer): **2.2**

Disturbance severity: **0.6**

Disturbance severity (herb layer): **0.6**

Mowing frequency: **1.09**

Grazing pressure: **0.22**

Soil disturbance: **0.16**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) Molinio-Arrhenatheretea](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R34 Submediterranean moist meadow](#)

Constant species of EUNIS habitats: [R34 Submediterranean moist meadow](#)

Dominant species of EUNIS habitats: [R34 Submediterranean moist meadow](#)

### Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Grassland (non-alpine, non-saline)**

## **Distribution**

Continentality: **5**

Continentality amplitude: **3**