

Alyssum turkestanicum

Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.46**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **5.8**

Moisture indicator value: **3.3**

Reaction indicator value: **8**

Nutrient indicator value: **1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.84**

Disturbance frequency (herb layer): **1.94**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.51**

Mowing frequency: **0.32**

Grazing pressure: **0.37**

Soil disturbance: **0.31**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CK \(SED\) Sedo-Scleranthetea](#), [DA \(FES\) Festuco-Brometea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1C Desert steppe](#), [S67 Aralo-Caspian semi-desert](#), [S68 Semi-desert sand dune with sparse scrub](#)

Constant species of EUNIS habitats: [R11 Pannonian and Pontic sandy steppe](#), [R1C Desert steppe](#), [S67 Aralo-Caspian semi-desert](#), [S68 Semi-desert sand dune with sparse scrub](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Semi-desert**