

Ambrosina bassii

Habitus and growth type

Plant height [m]: **0.05**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **January-March, November-December**

Fruit, seed and dispersal

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **7.5**

Moisture indicator value: **4**

Reaction indicator value: **4**

Nutrient indicator value: **4.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.28**

Disturbance frequency (herb layer): **1.43**

Disturbance severity: **0.68**

Disturbance severity (herb layer): **0.29**

Mowing frequency: **0.09**

Grazing pressure: **0.3**

Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **[FI \(TUB\) Helianthemetea guttati](#)**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**