Anabasis aphylla

Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Disturbance indicator values

Disturbance frequency: **2.35**
Disturbance frequency (herb layer): **2.2**
Disturbance severity: **0.35**
Disturbance severity (herb layer): **0.35**
Mowing frequency: **0.01**
Grazing pressure: **0.43**
Soil disturbance: **0.26**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **EA (LER) *Artemisietea lerchianae*, EC (KAL) *Kalidietea foliati***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S67 Aralo-Caspian semi-desert**
Constant species of EUNIS habitats: **S67 Aralo-Caspian semi-desert**

Broad habitat

Occurrence in broad habitats: **Saline vegetation, Semi-desert**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz