Anacamptis coriophora

Habitus and growth type

Plant height [m]: **0.22**
Life span: **Perennial**
Life form: **Geophyte**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **initial or partial mycoheterotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**
Reaction indicator value: **5.8**
Nutrient indicator value: **2.9**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.03**
Disturbance frequency (herb layer): **1.97**
Disturbance severity: **0.53**
Disturbance severity (herb layer): **0.47**
Mowing frequency: **0.74**
Grazing pressure: **0.31**
Soil disturbance: **0.22**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CM (MOL) *Molinio-Arrhenatheretea*, DA (FES) *Festuco-Brometea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

Distribution

Continentality: **6**
Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz