

# *Anagallis tenella*

## Habitus and growth type

Plant height [m]: **0.08**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **35.1**

## Flower

Flowering period: **May-July**

## Fruit, seed and dispersal

Seed mass [mg]: **0.11**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.7**

Moisture indicator value: **8.9**

Reaction indicator value: **4.8**

Nutrient indicator value: **2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.57**

Disturbance frequency (herb layer): **1.65**

Disturbance severity: **0.32**

Disturbance severity (herb layer): **0.31**

Mowing frequency: **0.33**

Grazing pressure: **0.29**

Soil disturbance: **0.2**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **OB (LIT) *Littorelletea uniflorae*, PA (SCH) *Scheuchzerio palustris*-*Caricetea fuscae***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **Ph Oligotrophic-water vegetation**

Constant species of EUNIS habitats: **Ph Oligotrophic-water vegetation**

### Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Mire**

## Distribution

Continentality: **2**

Continentality amplitude: **2**