

# *Androsace chamaejasme*

## Habitus and growth type

Plant height [m]: **0.03**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **May-July**

## Fruit, seed and dispersal

Seed mass [mg]: **0.52**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **2**

Moisture indicator value: **4.7**

Reaction indicator value: **8.3**

Nutrient indicator value: **2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.32**

Disturbance frequency (herb layer): **1.43**

Disturbance severity: **0.3**

Disturbance severity (herb layer): **0.29**

Mowing frequency: **0.02**

Grazing pressure: **0.25**

Soil disturbance: **0.08**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CU \(SES\) \*Elyno-Seslerietea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R44 Arctic-alpine calcareous grassland](#)

### Broad habitat

Occurrence in broad habitats: **Spring, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **7**

Continentality amplitude: **4**