Androsace obtusifolia

Habitus and growth type

Plant height [m]: **0.04**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.7**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Dystrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**
Temperature indicator value: **1.8**
Moisture indicator value: **5**
Reaction indicator value: **1**
Nutrient indicator value: **1**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.45**
Disturbance frequency (herb layer): **1.6**
Disturbance severity: **0.25**
Disturbance severity (herb layer): **0.24**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.02**
Grazing pressure: **0.23**
Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CT (TRI) *Juncetea trifidi***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz